

T: +27 (0) 35 791 5430
M: +27 (0) 82 956 3310 (emergency)
E: drlelala@ufhc.co.za
A: Suite No. 113
Melomed Private Hospital
John Ross Highway | Richards Bay, 3900

PLASTIC & RECONSTRUCTIVE SURGEON

[BSc, MBCHB, DA, PDBA, MMED (UCT), FC Plast Surg (SA)] | MP 0650269 | PR No. 0834815

Patient Information

Breast Augmentation

- It is the second most common Cosmetic surgery done after liposuction
- There are two types in implants used: Saline implants and Silicone implants
- Silicone implants have evolved through different generations and are currently approved for Breast augmentation and Breast reconstruction
- More details on implant types, textures, advantages and disadvantages of each will be discussed in detail during a consultation

Indications

- ☑ Enhance Breast shape and volume
- ☑ Improve body image, symmetry, and balance
- ☑ Helps clothes fit better
- ☑ Provide the appearance of a breast lift and increase cleavage
- ☑ Rejuvenation after postpartum deflation

Contraindications

- Patients with Body dysmorphic disorder
- **■** Patients under age of 18 years
- ☑ Significant breast disease & high risk of breast cancer
- **▼** Psychological instability
- Smokers (must stop smoking at least 3 weeks before the operation)



T: +27 (0) 35 791 5430
M: +27 (0) 82 956 3310 (emergency)
E: drlelala@ufhc.co.za
A: Suite No. 113
Melomed Private Hospital
John Ross Highway | Richards Bay, 3900

PLASTIC & RECONSTRUCTIVE SURGEON

[BSc, MBCHB, DA, PDBA, MMED (UCT), FC Plast Surg (SA)] | MP 0650269 | PR No. 0834815

Patients Evaluation

- > History
- **Examination**
- > Measurements




Fig. 2.5 Preoperative measures (taken before breast augmentation) include SSN:N (suprasternal notch to nipple); N:IMF (nipple to inframammary fold); BW (breast width); BH (breast height), and IMD (intermammary distance).

Incision Choices for introducing the Implant

- 1. Inframammary fold
- 2. Peri areolar
- 3. Trans axillary
- 4. Trans umbilical

Pocket plane & Dissection

- A. Sub glandular
- B. Subpectoral




T: +27 (0) 35 791 5430
M: +27 (0) 82 956 3310 (emergency)
E: drlelala@ufhc.co.za
A: Suite No. 113
Melomed Private Hospital
John Ross Highway | Richards Bay. 3900

PLASTIC & RECONSTRUCTIVE SURGEON

[BSc, MBCHB, DA, PDBA, MMED (UCT), FC Plast Surg (SA)] | MP 0650269 | PR No. 0834815

C. Biplanar (Dual Plane)

Pre & Post-op examples:



Potential Complications

- Capsular Contracture
- ❖ Implant leakage & rupture
- Capsular Calcification
- Asymmetry
- Altered nipple sensation
- **❖** Implant migration
- ❖ Anaplastic Large Cell Lymphoma (ALCL)



T: +27 (0) 35 791 5430
M: +27 (0) 82 956 3310 (emergency)
E: drlelala@ufhc.co.za
A: Suite No. 113
Melomed Private Hospital
John Ross Highway | Richards Bay, 3900

PLASTIC & RECONSTRUCTIVE SURGEON

[BSc, MBCHB, DA, PDBA, MMED (UCT), FC Plast Surg (SA)] | MP 0650269 | PR No. 0834815

Post-op Care

- ✓ Post op garment for 4 weeks, no underwire or push-up bra for 6 weeks
- ✓ Implant displacement exercise for 2 months
- ✓ Aerobic exercise after 2 weeks
- ✓ Heavy lifting after 6 weeks
- ✓ Back to work after 3 to 7 days

Disclaimer: for comprehensive medical information you can book a consultation appointment where an assessment will be made.